STANFORD ICHTHYOLOGICAL BULLETIN

VOLUME 8

Comprising 4 numbers, 8 articles,
259 pages, 73 figures, and 2 tables.

Published by the
DIVISION OF SYSTEMATIC BIOLOGY, STANFORD UNIVERSITY
Stanford, California, U.S.A.
1962-1967

IMPORTANT NOTICES

NO INDEX TO VOLUME 8 WILL BE ISSUED.

The Editors hereby affirm that copies of each of the four numbers of Volume 8 were mailed, on or before the imprinted date of publication, to the principal zoological, ichthyological, or natural history museum libraries of Ann Arbor, Berkeley, Cambridge, Chicago, Honolulu, Miami, New Haven, New York, Philadelphia, San Francisco, Seattle, Washington, Berlin, Calcutta, Cape Town, Copenhagen, La Plata, Leningrad, London, Paris, Rio de Janeiro, and Sydney, and to the editors of "Science" and "Nature." On the same day copies were placed on public sale at the Division of Systematic Biology, Stanford. The remainder of the copies distributed by the Division were mailed upon or within a few days of the publication date. Copies distributed through Stanford University Library were frequently delayed for a short time.

The Bulletin is issued irregularly, as publication funds come to hand. Because of this, and because of the lack of staff to handle them, we are unable to accept regular subscriptions. The Bulletin is offered in exchange for serials or other publications bearing upon ichthyology. A price list of back numbers is available, but many issues are now out-of-print and unavailable. In regard to purchase or exchange, please address the Director, Division of Systematic Biology, Stanford University, Stanford, California (94305), U.S.A.

Volume 8 was edited by G. S. Myers with the help of W. C. Freihofer and T. R. Roberts.

CONTENTS OF VOLUME 8

FREIHOFER, W. C. — Patterns of the Ramus Lateralis Accessorius and their systematic significance in teleostean fishes 81
——— New distributional records of the butterfly fish Chaetodon falcifer 207

MYERS, G. S., and W. C. FREIHOFER — Megalomycteridae, a previously unrecognized family of deep-sea cetomimiform fishes based on two new genera from the North Atlantic 193

MYERS, G. S., and T. R. ROBERTS — Note on the dentition of Creagrudite maxillaris, a characid fish from the Upper Orinoco-Upper Rio Negro system 248

ROBERTS, T. R. — Description and osteology of Lepidarchus adonis, a remarkable new characid fish from West Africa. . . . 209
——— Tooth formation and replacement in characoid fishes . . . 231
——— Virilia, a new genus of sexually dimorphic characid fishes from West Africa, with remarks on characoids having an incomplete lateral line. 251

WEITZMAN, S. H. — The osteology of Brycon meeki, a generalized characid fish, with an osteological definition of the family 3